

Screening of glyphosate herbicide degrading bacteria from soil

K.G.Maske

Department of Microbiology,
Rajarshi Shahu Mahavidyalaya, Latur (M.S.)

Abstract:

Glyphosate (N-phosphonomethylglycine) is a broad spectrum most widely used systemic herbicide worldwide. The present study was performed for isolation and characterization of bacteria having the efficiency to grow in presence of glyphosate. An enrichment culture technique was used to obtain bacterial strains from soil by using mineral salt medium containing 0.25 G/L to 2.0 G/L concentrations of the glyphosate as sole carbon or phosphorus source. Isolates having ability to grow at maximum concentration were selected and subcultured in mineral salt medium containing 1.25g/L glyphosate as sole carbon and phosphorous source. Total 75 isolates were obtained after enrichment and plating on nutrient agar plate. 45 isolates were found to gram negative rod shaped bacteria. 25 isolates were gram positive spore former bacteria and five bacterial isolates gram positive nonspore forming bacteria. Five isolates GL1, GL2, GL3, GL4, GL5 were selected on the basis of growth study. Bacterial strains were identified by using standard microbiological characteristics as *Pseudomonas aeuroginosa* (GL1), *Bacillus spp.*, *Serratia marsces*, *Pseudomonas aeuroginosa* (GL4) *Pseudomonas aeuroginosa* (GL5). Microbial degradation of glyphosate determined by spectrophotometric analysis of inorganic phosphate and thin layer chromatography. These bacterial strains possess potential to degrade glyphosate and could be used for bioremediation of glyphosate-contaminated sites.

Key words: Herbicides, glyphosate -Excel MERA 71, biodegradation, bioremediation, consortium

Introduction:

Pesticides are chemicals applied to control pests on agricultural crops. Herbicides are pesticides widely used in agriculture for weed control. They specially kill weeds, grasses that compete with commercial crops and its application has become indispensable for agriculture. Glyphosate is a systemic nonselective herbicide used worldwide (Franz et al., 1997, Woodburn, 2000). Only a small fraction of the chemicals available to the microorganisms leading to potential impact of residues of herbicides in soil and water on human animal and crop health (Pimentel, 1995). The surfactant polyoxyethylene amine (POEA) used in the formulation of glyphosate is more toxic as compared to glyphosate alone (Atkinson, 1985). A number of herbicides have been introduced as pre- or post-emergence weed killer (Ayansina and Oso, 2006). These herbicides release unwanted residues in soil, which are ecologically harmful (Haney et al., 2000; Derksen et al., 2002). Herbicides not only affect the target organisms, but also affect microbial communities important in soil activities (Perucci and Scarponi, 1994) and poses a risk to the entire ecological system (Kalia and Gupta, 2004). Microbes have key roles in organic matter transformations, nutrient cycling and degradation of organic pollutants, including pesticides (Beck et al., 2005). Bioremediation is attractive, less expensive and eco friendly method for degradation of chemicals from contaminated site (Alexander, 1999).

Glyphosate is organophosphate herbicide and account for half of the pesticide used world wide (Franz et al. 1997, Woodburn; 2000). Excel MERA 71, is a widely used glyphosate based formulation. It belongs organophosphorus group of herbicides. It is broad spectrum systemic nonselective and post emergent herbicide used in INDIA for controlling weeds in agriculture, forestry, urban areas, and even in aquatic bodies.

Glyphosate inhibit an enzyme 5-enol pyruyl shikimate-3-phosphate synthase, which is involved in the synthesis of aromatic amino acids-tyrosine, tryptophan and phenylalanine biosynthesis in plants. It also acts as competitive inhibitor of phosphonate pyruvate which is one of the precursor for biosynthesis of aromatic amino acids. Hence it is effective in growing plant. Glyphosate possess tendency of absorption of soil particles and organic matters Thus limits bio availability to microorganisms. In soil half life of Glyphosate vary from 3 days to 141 days (Andrea et al. 2003, Annsworth et al. 1993). Biodegradation is considered to be the most important of the transformation processes controlling its persistence in soil (Araujo et al., 2003).

Therefore present study was performed for the isolation and identification, characterization of glyphosate utilizing bacteria in soil. Capability of these of these isolates to degrade glyphosate herbicide confirmed by determination of inorganic acid phosphate and chromatographic analysis.

Materials and methods:

Chemicals and media used:

The isopropylamine salt of glyphosate known as Excel MERA 171 purchased from local supplier of Latur district, (M.S.) INDIA.

For the enrichment of glyphosate degrading bacteria mineral salt medium (MSM) containing glyphosate as sole carbon and phosphorous source was used. The composition of MSM (in g/L) was: (1.5), NaCl (0.5), NH_4SO_4 (2), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.2), CaCl_2 (0.01) and $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0.001). pH (7.0 to 7.2). Trace mineral solution 2 mL was added to MSM. Composition Trace mineral solution (in mg/L), NaCl (0.5), KCl (0.5) NH_4SO_4 (2), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.2) CaCl_2 (0.01) and $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0.001) $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, (0.020). Flasks of MSM (100mL) after sterilization were incorporated with 0.25G, 0.50G, 0.75G, 1.0 G, 1.25G, 1.50G, 2.0 G filter-sterilized (0.2 μm filter) glyphosate.

Collection of soil sample:

Soil samples were collected from two agriculture fields having previous history glyphosate application from Latur district. Soil samples were collected from depth of 0-15 cm in sterile petriplates. On arrival to laboratory stored in refrigerator.

Enrichment of glyphosate utilizing bacteria

Effect of different concentrations of glyphosate on total viable count of bacteria

About 1 g of soil sample added to 100 m L sterile physiological saline, after uniform mixing centrifuged at 3000 rpm for 5 min. 1 m L soil suspension inoculated to each flask containing 0.25G/L 0.50G/L, 0.75G/L 1.0 G/L, 1.25G/L, 1.50G/L, 2.0/L glyphosate as sole source of carbon and phosphorus. Flasks were incubated for 72 hr in shaking incubator at 150 rpm and 30°C.

After incubation 0.1 m L culture inoculated on nutrient agar plates. Plates were incubated at 30°C for 24 hr. After incubation growth was measured in terms of colony forming units per m L (cfu/ml).

Isolation of glyphosate degrading bacteria

For the isolation of bacteria that were survived highest concentration of MERA 171, from the flask containing 1.25G/L concentration of glyphosate 1.0 ml of culture was transferred to fresh MSM incorporated with 1.25G/L glyphosate. Flasks were incubated for 72 hr in shaking incubator at 150 rpm and 30°C. After incubation 0.1 m L culture was inoculated on the mineral salt agar plate containing 1.25 G/L Excel MERA 171 as a sole source of carbon and phosphorous. Plates were incubated for 48 hr. for growth of glyphosate utilizing bacteria. Macroscopic and microscopic characteristics were studied.

Colonies of glyphosate degrading bacteria were purified by sub culturing on nutrient agar plates .After incubation sub cultured on nutrient agar slant. After growth slants were stored at 4° C for further study.

Identification of glyphosate degrading bacteria

Isolates were subjected to morphological, biochemical studies including Gram staining , motility, endospore staining(Schaffer and Fulton's),capsule staining(Maneval's).Standard biochemical test including carbohydrate fermentation (glucose, lactose, sucrose, mannose), enzyme activities-oxidase , catalase , amylase , gelatinase, Indole formation, methyl red test, VP test, citrate utilization. The isolates were identified according to Bergey's Manual of Systematic Bacteriology.

Growth study of glyphosate utilizing bacteria

Inoculums of glyphosate utilizing bacterial cultures GL1, GL2, GL-3, GL-4, GL-5 were prepared by inoculation of pure culture of bacteria in to flask containing nutrient broth. Flasks were incubated at 30°C for 24 hr. After incubation culture were centrifuged at 5000 g for 5 min. Cell pellet was washed twice with sterile distilled water and resuspended in 2m L sterile distilled water cell density adjusted to 0.3 O.D. at 660 nm. 1.0 ml of culture was transferred to fresh MSM incorporated with 1.25G/L glyphosate. Flasks were incubated for 15 days in shaking incubator at 150 rpm and 30°C. Growth was monitored in terms of cfu /m L at the interval of 24 hr. by inoculating 0.1 m L culture on nutrient agar plate and after incubation for 24hr at 30°C.

Determination of glyphosate biodegradation

Spectrophotometric analysis of inorganic phosphate

Culture (3mL) from the above flask were centrifuged at 5000 g for 5 min and supernatants were used for analysis of inorganic phosphate(Fiske and Subbaraw, 1925). To the 1.0 m L of the supernatant, 2mL Trichloroacetic acid, 1.0 m L ammonium molybdate solution and 1,2,4-aminonaphtholsulphonic acid was added. The mixture was thoroughly mixed and allowed to react for 5 minutes. The absorbance was read 660 nm using spectrophotometer

(Shimadzu) against blank. Analysis was carried out in triplicate for supernatant obtained from each flask at the interval of 24 hr. Control flasks (un inoculated) were also analyzed.

Thin layer chromatographic analysis

Cultures from the above flasks after incubation for 15 days were centrifuged at 5000 g for 5 min. Supernatant was extracted with acetonitrile and thin layer chromatography was carried out on TLC sheets pre coated with silica gel (Merck). Chromatography was performed using solvent system butanol-acetic acid-water (24:6:10). Spots were developed with ninhydrin reagent. Standard compounds glyphosate, glycine and sarcosine were used as the reference for identification of compounds.

Results and discussion

Enrichment and isolation of glyphosate utilizing bacteria

In order to obtain population of bacteria having ability to grow at higher concentration of glyphosate enrichment was carried out in different flasks containing mineral salt medium incorporated with 0.25G/L to 2.0G/L concentrations. Growth was monitored at the interval of 24 hr. in terms of cfu/ mL. Total count was decreased as concentration of glyphosate was increased. Marked decrease in cfu/m L was found at concentration 1.25G/L.




Fig.1: Total viable count of glyphosate utilizing bacteria at different concentrations of glyphosate

Initially total count of glyphosate utilizing bacteria was constant at the 0.25G/L and 0.50 G/L. Total count was decreased with increasing concentration up to 2G/L. Total 75 isolates were obtained after enrichment and plating of culture(1.25G/L) nutrient agar plate. 45 isolates were found to be gram negative rod shaped bacteria. 25 isolates were gram positive spore former bacteria and five bacterial isolates were gram positive nonspore forming bacteria. Only five isolates were predominated at 2mg/L glyphosate. Isolates were designated as GL-1, GL-2, GL-3, GL-4, GL-5. Glyphosate is an organophosphonate that can be used as a source of P, C or N by either gram positive or gram negative bacteria reported by van Eerd et al., 2003. Glyphosate was found to be lethal to bacteria above 50 mg/ L concentration. Glyphosate reduced bacterial viability on solid media and complete inhibition of fungal growth reported by Busse et al., 2000.

Identification of glyphosate degrading bacteria

Cultural, morphological and biochemical characters of five glyphosate degrading isolates were studied and isolates GL-1, GL-2, GL-3, GL-4, GL-5 were identified on the basis of Bergey's Manual of systematic Bacteriology. Table-1 and table-2

Table-1. Cultural characteristics of glyphosate degrading bacterial isolates

Isolate	Colony size	Shape	Chromogen sis	Surface	Elevation	Edge	Opacit y	Consistency
GL-1	2mm	circular	Yellowish	smooth	convex	entire	Translucent	Butyrous
GL-2	1mm	circular	white	smooth	flat	irregular	opaque	Butyrous
GL-3	2mm	circular	pink	smooth	umbonate	entire	Translucent	Butyrous
GL-4	1mm	circular	yellowish	smooth	raised	entire	Translucent	Butyrous
GL-5	3mm	circular	orange	smooth	convex	entire	Translucent	Butyrous

Table-1. Morphological and biochemical characteristics of glyphosate degrading bacterial isolates

Isolates	GL1	GL2	GL3	GL4	GL5
Characteristics					
Morphology	rod	rod	rod	rod	Rod
Gram staining	-	-	-	-	-
Endospore staining	-	+	-	-	-
Motility	+	+	+	+	+
Oxidase	+	+	-	+	+
Catalase activity	+	+	+	+	+
Indole production	-	-	-	+	-
MR	-	-	-	-	-
VP	-	+	+	-	-
Citrate utilization	+	+	+	+	+
Starch hydrolysis	-	+	-	-	-
Gelatinase	+	+	+	+	+
Sugar fermentation					
1.Glucose	-	+	-	-	-
2.Fructose	-	+	+	+	+
3.Sucrose	+	+	+	+	+
4.Lactose	-	+	+	+	-
5.Mannose	+		+	+	+
6.Mannitol	+		+	+	-
Arabinose	+	+	-	ND	ND
H ₂ S production	-		-	-	-
Lysine decarboxylation	+	+	+	+	-
Tentative identification	<i>Pseudomonas aeruginosa</i> (stra	<i>Bacillus pumilus</i>	<i>Serratia marcesce</i>	<i>Pseudomonas aeruginosa</i> (stra	<i>Pseudomonas aeruginosa</i> (stra

	in-1)		ns	in-2)	in-3)
--	-------	--	----	-------	-------

ND: Means not determined ,+ means positive test,- means negative test

Five bacterial isolates GL1, GL2, GL3, GL4, GL5 were tentatively identified as *Pseudomonas aeuroginosa*(strain-1), *Bacillus pumilus* , *Serratia marcescens*, *Pseudomonas aeuroginosa*(strain -2), *Pseudomonas aeuroginosa*(strain-3)

Growth study of glyphosate degrading bacteria

Growth study of glyphosate degrading bacteria carried out by using mineral salt medium containing 2G/L glyphosate. Strain GL1, GL2, GL3 showed maximum growth 4.2 Log 10 cfu/ml, 3.2 Log 10 cfu/ml, 3.7 Log 10 cfu/ml after twelve days of incubation. Strain GL4, GL5 showed 3.9 Log 10 cfu/ml , 4.2 Log 10 cfu/ml after incubation for 10 days. Growth kinetics of glyphosate degrading bacteria was as shown in figure 2a and figure 2b.

Spectrophotometric analysis of inorganic phosphate-

Glyphosate degradation by isolated bacteria was determined by monitoring spectrophotometric analysis of inorganic phosphate using Fiske Subbarow method Concentration of inorganic phosphate determined at the interval of 24 hr in the culture broth containing 2G/L of glyphosate. Results were as per table-3

Chromatographic analysis of glyphosate biodegradation:

Chromatographic analysis of extracted biodegradation products from both *Pseudomonas aeuroginosa* GL4 showed sarcosine Rf 0.54 Chromatographic analysis of extracted biodegradation products from culture of *Bacillus pumilus*. showed presence of sarcosine (Rf=0.54) and glycine Rf. 0.25 .Complete degradation of glyphosate observed after 15 days by GL4-*Pseudomonas aeuroginosa* , *Serratia marcescences*- showed presence of glycine Rf-0.25

Discussion :

Glyphosate is an organophosphonate that can be used as a source of P,C or N by either gram positive or gram negative bacteria reported by van Eerd et al.,2003.Application of herbicide to the soil may adversely affect sensitive microorganisms and simultaneously those microorganisms that have ability to degrade herbicide get stimulated.(Wikinson and Lucas,1969;Ayansina and Oso,2006) Glyphosate was found to be lethal to bacteria above 50 mg/ L concentration and glyphosate reduced bacterial viability on solid media and complete inhibition of fungal growth reported by Busse et al.2000.In the present study also marked inhibition observed at higher concentration above 200mg/L concentration. However ,Amoros et al.2007,studied effect of different concentration of glyphosate on Aeromonas count and reported increase in count as compared to the control. M. Baboo et al. (2013), reported that herbicide treatment resulted an initial increase in biomass upto 14th day followed by significant drop in biomass and indicated transient impact herbicide.

Capacity to metabolize of glyphosate have been reported in Several species of *Pseudomonas* (Moore et al., 1983; Jacob et al., 1988; Quinn et al., 1989). Metabolism of glyphosate by *Bacillus megaterium* strain 2BLW (Quinn et al., 1989) also reported , *B. cerus* (Fang et al.2012).

Three *Pseudomonas aeuroginosa* strains, *Bacillus pumilus* , *Serratia marcescens* were found to be predominated while other population was found to be inhibited at 2G/L. Growth study of

all the *Pseudomonas aeuroginosa* strains showed higher growth as compared to *B. pumilus* and *Serratia marcescens*. These observations are consistent with studies of Gimsing et al.(2004) who reported a high correlation between glyphosate mineralization rate and *Pseudomonas* sp counts in five different Danish soils.

Microbial degradation of glyphosate produces the major metabolite aminomethylphosphonic acid and ultimately leads to the production of CO₂, phosphate and water (Forlani et al.,1999;Araujo et al.,2003) In the present study concentration of inorganic phosphate determined at the interval of 24 hr in the culture broth containing 2G/L of glyphosate. Concentration of inorganic phosphate in the culture of *Pseudomonas aeuroginosa* found to be increased during 144 hr of incubation it indicate increased transformation of glyphosate. In case of *Bacillus pumilus*, *Serratia marcescens* no significant concentration was detected, it might be utilized for growth.




Figure2a. Growth study of glyphosate degrading bacteria in medium containing glyphosate as sole source of carbon and phosphorous




Figure2b. Growth study *Pseudomonas aeuroginosa* (GL1) and *Pseudomonas aeuroginosa* (GL5) in medium containing glyphosate as sole source of carbon and phosphorous




Figure 3: Analysis of inorganic phosphate in broth cultures containing glyphosate

Chromatographic analysis of glyphosate biodegradation:

Chromatographic analysis of extracted biodegradation products from both *Pseudomonas aeuroginosa* showed presence of AMPA. Chromatographic analysis of extracted biodegradation products from culture of *Bacillus pumilus* showed presence of degradation product sarcosine and glycine, *Serratia marcescens* showed presence of glycine.

Pseudomonas sp. PG2982 degrade glyphosate by initial cleavage of the C-P bond resulting in the production of sarcosine(N-methyl-glycine) by C-P lyase activity ((Kishore & Jacob, 1987).

Conclusion

Present study confirmed ability of *Pseudomonas aeuroginosa* strains, *Bacillus pumilus*, *Serratia marcescens* to degrade glyphosate. Chromatographic analysis and analysis of inorganic phosphate confirmed transformation of glyphosate in vitro by these isolates. Consortium of these microorganisms may be employed for detoxification of contaminated sites.




Fig. Thin layer chromatography

1. Complete degradation of glyphosate after 15 days by GL4-*Pseudomonas aeuroginosa*,
2. *Serratia marcescences*-Presence of glycine, Rf-0.25
3. *Bacillus pumilus*--Presence of glycine, Rf. 0.25 and sarcosine Rf 0.54

4.glycine Rf. 0.25

S-Glyphosate Rf 0.12

GL4-*Pseudomonas aeuroginosa* -Sarcosine Rf 0.54

References

- A. Kalia, R.P. Gupta, Disruption of food web by pesticides, *Ind. J. Ecol.* 31 (2004) 85-92.
- A.D.V. Ayansina, B.A. Oso, Effect of two commonly used herbicides on soil microflora at two different concentrations, *Afri. J. Biotech.* 5(2) (2006) 129132. 7
- Ainsworth, C. C., Frederickson, J. K. & Smith, S. C. (1993). Effect of sorption on the degradation of aromatic acids and bases, in D. M. Linn, T. H. Carski, M. L. Brusseau & F. H. Chang (eds), *Sorption and Degradation of Pesticides and Organic Chemicals in Soil*, Soil Science Society of America, Madison, WI, pp. 125–144.
- Alexander M. Biodegradation and bioremediation, San Diego, USA; Academic Press, ISBN 978-0-12-049861-1, 1999, pp 453.
- Andrea MM, Tomita RY, Luchini LC, Musumeci MR, 1994. Laboratory studies on volatilization and mineralization of 14c-p,p-DDT in soil, release of bound residues and dissipation from solid surfaces; *Journal of Environmental Science and Health, Part B*, Volume 29, Issue 1, 1994, pp 133-139.
- Araujo ASF, Monteiro RTR & Abarkeli RB (2003) Effect of glyphosate on the microbial activity of two Brazilian soils. *Chemosphere* 52: 799–804.
- Atkinson D (1985). Toxicological properties of Glyphosate-A Summary. In : *The Herbicide glyphosate*. E. Grossbard and D. Atkinson (eds.). Butterworths, London, U.K. p.127.
- Balthazor TM, Hallas LE (1986). Glyphosate degrading microorganisms from industrial activated sludge. *Appl. Environ. Microbiol.* 51:432-434
- Beck, L., Römbke, J., Breure, A. M. & Mulder, C. (2005). Considerations for the use of soil ecological classification and assessment concepts in soil protection, *Ecotoxicology and Environmental Safety* 62: 189–200.
- Benoit, P., Barriuso, E. & Soulas, G. (1999). Degradation of 2,4-D, 2,4-dichlorophenol and 4-chlorophenol in soil after sorption on humified and nonhumified organic matter, *Journal of Environmental Quality* 28: 1127–1135.
- Busse, M. D., Ratcliff, A. W., Shestak, C. J. & Powers, R. F. (2000). Non target effects of glyphosate on soil microbes, *Proceedings of the California Weed Science Society* 52:146-150.
- Derksen D.A., Anderson R.L., Blackshaw R.E. and Maxwell B., Weed dynamics and management strategies for cropping systems in the Northern Great Plains, *Agron. J.* 94 (2002) 174–185.
- Fan J, Yang G, Zhao H, Shi G, Geng Y, Hou T, Tao K. (2012). Isolation, identification, and characterization of a glyphosate-degrading bacterium, *Bacillus cereus* CB4, from soil. *J. Gen Appl Microbiol.* 58(4):263-71
- Franz JE, Mao MK, Sikorski JA 1997. *Glyphosate: A unique global herbicide*. American Chemical Society Monograph 189. American Chemical Society, Washington DC.
- Jacob GS, Gabrow JR, Hallas LE, Kimack NM, Kishore GM, Schaefer J (1988). Metabolism of Glyphosate in *Pseudomonas* sp. Strain LBr. *Appl. Environ. Microbiol.* 54:2953-2958.
- Kishore, G. M. & Jacob, G. S. 1987. Degradation of glyphosate by *Pseudomonas* sp. PG2982 via a sarcosine intermediate, *Journal of Biological Chemistry* 262: 12164–12168.
- Kitagawa van Eerd, L. L., Hoagland, R. E., Zablotowicz, R. M. & Hall, J. C. (2003). Pesticide metabolism in plants and microorganisms, *Weed Science* 51: 472–495

- Riaz M., Jamil M., Mahmood T.Z., Yield and yield components of maize as affected by various weed control methods under rain-fed conditions of Pakistan, *Int. J. Agric. Biol.* 9 (2007)152–155.
- Moore IK, Braymer HD & Larson AD (1983) Isolation of a *Pseudomonas* sp. which utilises the phosphonate Perucci P., L. Scarponi, Effects of the herbicide imazethapyr on soil microbial biomass and various soil enzyme activities, *Biol. Fertility and Soils.* 17 (1994) 237-240.
- Pimentel, D. (1995). Amounts of pesticides reaching target pests: Environmental impacts and ethics, *Journal of Agricultural and Environmental Ethics* 8: 17–29.
- Quinn JP, Peden JMM & Dick RE (1989) Carbon-phosphorus bond cleavage by gram-positive and gram-negative soil bacteria. *Appl Microbiol Biotechnol* 31: 283–287.
- Pipke R and Amrhein N.(1988) . Degradation of the Phosphonate Herbicide Glyphosate by *Arthrobacter atrocyaneus* ATCC 13752. *Applied and environmental microbiology*, p. 1293-1296.
- Haney R. L., Senseman S. A., Hons F. M., Zuberer D.A., Effect of glyphosate on soil microbial activity and biomass, *Weed Sci.* 48(2000)89-93.
- Wilkinson V., Lucas R. L., Effects of herbicides on the growth of soil fungi, *New Phytol.* 68(3) (1969) 709-719.
- Woodburn,A.T.(2000).Glyphosate:production, pricing, and use worldwide, *Pest Management Science* 56:309-312.
- Baboo M, Mamata Pasayat, Alka Samal, Monty Kujur, Jitesh Kumar Maharana and Amiya Kumar Patel (2013).Effect of four herbicides on soil organic carbon, microbial biomass-c, enzyme activity and microbial populations in agricultural soil,*International Journal of Research in Environmental Science and technology*3(4):100-112
- Amoros I,Alonso JL,Romaguera S, Carrasco JM2007.Assessment of toxicity of glyphosate-based formulation using bacterial systems in water.*Chemosphere*.67:2221-2228